



The Influence of Localized Winds on Arctic Freshwater and Heat Export

By Amber M. Holdsworth*, Paul G. Myers**



** University of Alberta,
Canada



-Special thanks to Xianmin Hu for technical support.
-We are grateful to Gregory Smith for the CGRF forcing data.
-Thanks to Laura de Steur for the Fram Strait Transports.
-Many undergraduate students also contributed in some way to this project: Zoe Cire, Sherry Gao, David Stanfel, Jie Jian, Christina Chan.

Motivation

- Bursts of intense wind can enhance air-sea heat exchange
-> Trigger deep or intermediate convection [1]
that impacts:
 1. nutrient distribution and chemical exchange [2]
 2. Atlantic Meridional Overturning Circulation (AMOC) [3]
- as the climate warms storms are predicted to increase [4]
- How do bursting winds affect the transport of heat and freshwater out of the Arctic?

Bursting winds:
storms, barrier winds, cold air outbreaks,
polar lows, topographic jets

Ocean-Ice Model

- NEMO -Nucleus for European Modeling of the Ocean [5]
- Arctic Northern Hemisphere Atlantic Configuration
- LIM2 - Louvain-la-Neuve Sea Ice Model (LIM)
- 1/4° grid
- 50 vertical levels

Atmospheric Forcing

- CGRF-Canadian Meteorological Centre's Global Deterministic Prediction System [7]
- hourly atmospheric data (2002-2014)
- Resolution: 0.45° longitude, 0.3° latitude
- model forecasts from operational analysis




Fig.1: Model configuration and bathymetry.

Transport Validation




Fig.2: Locations for mooring array (pink) and regional mask (orange).

Correlation	Mean Values
0.47	data: -8.16 Sv model: -8.056 Sv

Fig.3: Monthly averaged volume transports from a mooring array in the East Greenland Current [8] plotted with the equivalent section in the model.

Filtering the Atmospheric Forcing

- wind and temperature fields were filtered using a moving average filter [6]
- time scales less than one week were removed




Fig.4: Wind stress.




Fig.5: Wavelet spectrogram of filtered and unfiltered data.

Bursting Winds Influence Arctic Transports




Fig.6: Transport sections (orange and purple (shown)) and the Arctic region used to compute Heat content (yellow).




Fig.7: Transports taken through the purple sections in Fig.6. Transports out of the Arctic are positive.

Bursting winds:

- enhance southward transport of freshwater
- play a role in heat transport into the Arctic

Heat Content in the Arctic

1. Over the Entire Depth




Fig.8: Averages taken within the Arctic basin (yellow curve in Fig.6).

Bursting winds:

1. • decrease the FW content
• lead to a colder Arctic Basin
2. • deepen and warm the mixed layer
• over time the difference between the two simulations increases

2. In the Mixed Layer




Fig.9: Averages taken within the Arctic basin (yellow curve in Fig.6). We normalize by the depth of the mixed layer.

Fram Strait

What was removed by the filtering?




Fig.10: Averaged over the orange rectangle in Fig.3

Filtered vs unfiltered




Fig.11: Transports taken across the yellow line marked 4 in Fig. 6. Positive indicates southward flow.

Bursting winds:

- inhibit transport of freshwater out of the Arctic
- increases the southward flow of heat

Directional Constancy




Fig. 12 Winter climatology of the CGRF 10 m winds North of Greenland from 2002 - 2010 (DJF). With sea level indicated in black the color-map shows the directional constancy (DC) of the wind, dark contours are labelled with the scalar averaged wind speeds, and mean wind vectors are plotted on top.

A directional constancy near one implies that the wind tends to flow in a single direction

$$DC = \frac{\bar{S}_u}{\bar{S}} = \frac{\sqrt{\bar{u}^2 + \bar{v}^2}}{\sum_{i=1}^N \sqrt{u_i^2 + v_i^2}}$$

- the wind tends to flow southward
- northward moving storms may affect transports
- strong topographically accelerated flows enhance the downward heat flux.

Gale Force Wind Events in Fram Strait

Criterion for an event:

- events begin when wind speeds exceed 17 m/s for at least 6 hours anywhere in the study region shown in orange in Fig.3.
- An event ends when winds are below 17m/s for at least 3 hrs
- there are 24 hrs between events

- Volume and freshwater transports were computed through the yellow section in Fig. 2.

- We found the relative difference between the transports occurring during an event and the average over the month in which the event occurred.

Gale force wind events can strongly enhance transports and reverse the direction of flow




Fig.13: red markers indicate that the direction of transport during the event is opposite to the monthly average.

Summary

Bursting winds ...

- significantly impact the transport of freshwater out of the Arctic
- Influence the heat content of the Arctic basin

Future directions and unanswered questions:

- Bursting winds bring warmer waters northward and mix warmer surface waters leading to a warmer mixed layer, but why/how do they cool the Arctic basin?
 1. separate the impacts of different winds (cyclones vs. tip jets etc.)
 2. analysis of transports at various depths
 3. examine ssh and dynamic heights for each of the straits
 4. examine FW pathways
- In the CAA, the Eurasian basin and around Baffin bay bursting winds enhance FW exports, but these winds can inhibit FW export through Fram Strait.
 1. separate the impacts of different winds (cyclones vs. tip jets etc.)
 - >self-organized maps
 2. analysis of transports at various depths

References

- [1] Holdsworth, Amber M., and Paul G. Myers. "The Influence of high-frequency atmospheric forcing on the circulation and deep convection of the Labrador Sea." *Journal of Climate* 28.12 (2015): 4980-4996.
- [2] Azetsu-Scott, K., Jones, E. P., Yashayev, I., & Gershney, R. M. (2003). Time series study of CFC concentrations in the Labrador Sea during deep and shallow convection regimes (1991-2000). *Journal of Geophysical Research: Oceans* (1978-2012), 108 (C11).
- [3] Kuhlbrodt, T., Griesel, A., Montoya, M., Levermann, A., Hofmann, M., & Rahmstorf, S. (2007). On the driving processes of the Atlantic Meridional Overturning Circulation. *Reviews of Geophysics*, 45 (2).
- [4] Zahn, M., & von Storch, H. (2010). Decreased frequency of North Atlantic polar lows associated with future climate warming. *Nature*, 467 (7313), 309-312.
- [5] Madec, G. (2008). NEMO reference manual, ocean dynamic component. NEMO-OPA. Note du Pôle de modélisation, Institut Pierre Simon Laplace, Technical Report, 27.
- [6] Eskridge, R. E., Ku, J. Y., Rao, S. T., Porter, P. S., & Zurbenko, I. G. (1997). Separating different scales of motion in time series of meteorological variables. *Bulletin of the American Meteorological Society*, 78(7), 1473-1483.
- [7] Smith, G. C., Roy, F., Mann, P., Dupont, F., Brasnett, B., Lemieux, J. F., ... & Bélair, S. (2014). A new atmospheric dataset for forcing ice-ocean models: Evaluation of reforecasts using the Canadian global deterministic prediction system. *Quarterly Journal of the Royal Meteorological Society*, 140 (680), 861-894.
- [8] de Steur, L., et al. "Impact of recirculation on the East Greenland Current in Fram Strait: Results from moored current meter measurements between 1997 and 2009." *Deep Sea Research Part I: Oceanographic Research Papers* 92 (2014): 26-40.